

INSTRUCTIONS FOR OUTPATIENT PROCEDURES/COLONSCOPY

HALF GOLYTELY WITH BISACODYL (DULCOLAX) PREPARATION

- 1. **FIVE DAYS PRIOR TO PROCEDURE: (Restricted Residue Diet) DO NOT EAT** nuts, seeds, popcorn, or corn. Discontinue fiber supplements such as Metamucil, Citrucel, Fiberall, etc.
- 2. ONE DAY PRIOR TO PROCEDURE: You may only eat or drink clear liquids the entire day prior to your test. You may have Jell-O, popsicles, clear juices that have no pulp, black coffee, tea, colas, etc. NOTHING RED, PURPLE, OR BLUE AND NO DAIRY PRODUCTS!!!
- 3. At 12:00 p.m. (noon) the day prior to the procedure, you should take 2 bisacodyl tablets (Dulcolax) with water.
- 4. Add lukewarm water to the top of the line on bottle and shake to dissolve powder.
- 5. At 6:00 p.m. the day before your test, begin your Golytely solution.
- 6. Drink 1 (8oz) glass of solution every 20-25 minutes until you have consumed 2 liters (1/2) of the solution. This solution should give you diarrhea so that your colon is clean for a good examination at the time of your test. If you do not drink all of the solution, your examination will be inadequate and may need to be rescheduled.
- 7. On the day of your procedure do not drink anything two hours prior to your arrival time. It is okay to take your required medication with only a small amount of water.