

INSTRUCTIONS FOR OUTPATIENT PROCEDURES/COLONSCOPY

HALFLYTELY PREPARATION

- FIVE DAYS PRIOR TO PROCEDURE: (Restricted Residue Diet) DO NOT EAT nuts, seeds, popcorn, or corn. Discontinue fiber supplements such as Metamucil, Citrucel, Fiberall, etc.
- 2. **ONE DAY PRIOR TO PROCEDURE:** You may only eat or drink clear liquids the entire day prior to your test. You may have Jell-O, popsicles, clear juices that have no pulp, black coffee, tea, colas, etc. **NOTHING RED, PURPLE, OR BLUE AND NO DAIRY PRODUCTS!!!**
- 3. At 12:00 p.m. (noon) the day prior to the procedure, you should take 2 bisacodyl tablets with water.
- 4. Add lukewarm water to the top of the line on bottle and shake to dissolve powder.
- 5. After a bowel movement occurs (usually in 1 to 6 hours) begin drinking the solution (even if no bowel movement occurs after 6 hours, begin to drink solution).
- 6. Drink 1 (8oz) glass of solution every 10-15 minutes until all of the solution is gone. This solution should give you diarrhea so that your colon will be cleaned for a good examination at the time of your test. If you do not drink all of the solution, your examination will be inadequate and may need to be rescheduled.
- 7. On the day of your procedure do not drink anything two hours prior to your arrival time. It is okay to take your required medication with only a small amount of water.